

Large N QCD on the lattice – A review of recent results in the fermionic sector

R. Narayanan

*Department of Physics,
Florida International University,
Miami, FL 33199, USA
E-mail: narayanr@fiu.edu*

H. Neuberger

*Department of Physics and Astronomy,
Rutgers University,
Piscataway, NJ 08855, USA
E-mail: neuberger@physics.rutgers.edu*

Abstract

It is possible to numerically solve QCD in the planar limit using standard numerical techniques on existing computer clusters. The basic ideas behind the computational strategy and recent numerical results in the fermionic sector of large N QCD are reviewed.

I. INTRODUCTION

The large N limit of QCD has several attractive features:

- Only planar diagrams contribute at fixed g^2N [1].
- Fermions in the fundamental representation are automatically quenched as long as the number of flavors is kept finite when taking $N \rightarrow \infty$ [1].
- It is expected to preserve the essential perturbative and non-perturbative features of the $N = 3$ theory. The exact solution of the two dimensional theory reveals a tower of mesons [2].
- Phenomenological predictions from large N QCD tend to be close to experimental results indicating that leading corrections in the $1/N$ expansion are at least occasionally small [3].
- Recent work indicates a duality between large N gauge theory and some string theory at zero string coupling [4].

Lattice QCD provides a non-perturbative method to numerically solve large N QCD. Conventional methods involve computing physical quantities as a function of N and extrapolating these results to the $N \rightarrow \infty$ limit [5]. So far, this approach has only been used to compute observables in the pure gauge theory, but it has provided evidence that large N can be quantitatively useful even at $N = 3$. Conventional computations proceed as follows: One first takes the large volume limit followed by the continuum limit at a fixed N and then extrapolates to $N \rightarrow \infty$. This approach will be difficult for fermionic quantities. This is due to the fact that the quenched theory at finite N has pathological singularities [6]. Therefore, it is necessary to first compute fermionic observables in a full dynamical simulation at a fixed N and then take the $N \rightarrow \infty$ limit.

A short cut to obtain results in the $N \rightarrow \infty$ limit using lattice QCD should use two aspects of large N QCD. One is that the physical volume corrections are expected to be small in the $N \rightarrow \infty$ limit since one has infinite number of degrees of freedom at a single lattice site. Secondly, fermions are naturally quenched in the $N \rightarrow \infty$ limit. To make use of the second feature, it is best to take the $N \rightarrow \infty$ limit of a quenched theory at a fixed lattice coupling, $b = 1/(g^2N)$, before taking the large volume limit and the continuum limit.

Since we are in a fixed lattice volume and lattice coupling, all fermionic observables are well-defined and therefore the $N \rightarrow \infty$ limit will be well-defined. Since we are taking the large volume limit in the $N \rightarrow \infty$ theory, this is also well-defined and does not suffer from quenched pathologies. Lastly, we are taking the continuum limit of a set of well-defined theories on the lattice.

The procedure described in the previous paragraph will work independent of the size of the finite volume corrections in the $N \rightarrow \infty$ limit. But, as mentioned before, this correction is expected to be small. In fact, tremendous excitement was generated by a paper of Eguchi and Kawai [7] in the early 1980s since it showed that the lattice theory in the $N \rightarrow \infty$ limit reduced to a single site model if the $U^d(1)$ symmetries specific to this lattice model remain unbroken. This theory has only d $SU(N)$ matrices with $N \rightarrow \infty$ and the $U^d(1)$ symmetries are the $N \rightarrow \infty$ limit of the Z_N^d symmetries associated with the d Polyakov loops in the single site model. Unfortunately, the $U^d(1)$ symmetries are broken in the weak coupling phase for $d > 2$ [8]. Since the reduction to a single site model is very attractive, solutions to fix the above problem were immediately found. One is referred to as the quenched Eguchi-Kawai model [8] and another is referred to as the twisted Eguchi-Kawai model [9]. The twisted Eguchi-Kawai model is an elegant solution to the problem encountered by the original proposal of Eguchi and Kawai and has also led to the discovery of non-commutative gauge field theories [10]. But, it is possible to include fermions only under some very specific conditions in the twisted Eguchi-Kawai model.

Quenched Eguchi-Kawai solves the problem of the original Eguchi-Kawai in an intuitive fashion and has other implications. The idea is to restrict the integral over all $SU(N)$ matrices with a fixed set of N eigenvalues. Observables are first computed under this restriction and these observables are then averaged over all choices for the N eigenvalues. Freezing the eigenvalues of the $SU(N)$ matrix in the computation of physical observables can be interpreted as quenched momentum prescription for gluons shown to be valid in the realm of perturbation theory [11]. In fact, the interpretation of quenched momentum prescription extends to fermions [12]. Fermion propagators are computed in the pure gauge theory owing to the fact that fermion loops are quenched in the $N \rightarrow \infty$ limit. But, fermion propagators with momenta that differ by p_μ in a fixed gauge field background are computed by using two different gauge field backgrounds that differ by a $U(1)$ phase of e^{ip_μ} in the d directions. This is referred to as the quenched momentum prescription for fermions. Since we integrate

over all $SU(N)$ fields, momentum for a single propagator has no meaning. Only physical observables like meson propagators carry momentum where the two quarks that make up the meson see gauge fields that differ by the $U(1)$ phase. Furthermore, one sees that mesons can have momentum as small as $2\pi/N$ on a single site lattice and therefore one can study mesons for a continuum of momenta on a single site lattice in the $N \rightarrow \infty$ limit.

The proposal to numerically solve QCD in the large N limit should therefore proceed as follows. Consider the single site $SU(N)$ lattice gauge theory with the Wilson plaquette action. This model has d $SU(N)$ matrices denoted by U_μ , $\mu = 1, \dots, d$. Generate gauge fields U according to the single site Wilson plaquette action with the added restriction that the d $SU(N)$ integrals go over all $SU(N)$ matrices with a fixed set of N eigenvalues that are uniformly distributed on the unit circle. Compute quark propagators in the fixed gauge field background $e^{ip_\mu} U_\mu$ where p_μ is restricted to take values equal to $\frac{2\pi n_\mu}{N}$ such that modified gauge field seen by the quark belongs to $SU(N)$. This quark propagator, $G(p, U)$ carries a momentum equal to p_μ . A meson propagator is computed using $\text{Tr} G(p, U) G(q, U)$ and the result is a meson carrying a momentum $P = p - q$. Averaging over all gauge field configurations results in the meson propagator in the large N limit of QCD. To recover “translational invariance”, it is necessary to sum over all p and q for a fixed $P = p - q$. This amounts to a computation of N^d propagators in a fixed gauge field background and this is a time consuming numerical computation for $d = 4$. But the idea of momentum quenching can be tested in $d = 2$ and the results [13] agree with the exact solution of ’t Hooft [2].

Eigenvalues of U_μ are uniformly distributed on the unit circle in the quenched Eguchi-Kawai model. Then, $e^{ip_\mu} U_\mu$ with $p_\mu = \frac{2\pi n_\mu}{N}$ is just a cyclic permutation of all the eigenvalues of U_μ and should be included in the average over all gauge fields. But the numerical algorithm that generates the gauge fields does not succeed in doing this and it is therefore necessary to sum over all p and q for a fixed $P = p - q$ when computing meson propagators. Since it is necessary to take the large N limit first, the proposal in the previous paragraph is quite a bit more numerical involved than one naively envisions. Therefore, it calls for a different realization of the original Eguchi-Kawai idea. The reduction argument of Eguchi and Kawai also applies to the lattice model on an L^4 periodic lattice. Reduction will go through and there will not be any finite volume effects if the d $U(1)$ symmetries associated with the d Polyakov loops are not broken. Indeed, numerical analysis shows that the above symmetries are not broken as long as $L > L_c(b)$ for $d = 3$ [14] and $d = 4$ [15]. Furthermore, $L_c(b)$

scales properly as one takes the continuum limit indicating the existence of a physical scale l_c such that large N QCD is properly reproduced, without finite volume effects if $l > l_c$ [15]. This is referred to as continuum reduction and it results in a modification of the proposal in the previous paragraph. One works on an L^d lattice with L just above $L_c(b)$ instead of working on a 1^d lattice. The gauge fields are generated using the standard single plaquette Wilson action with no restriction on the gauge fields themselves. All other aspects of the proposal remains the same, with the fermion momenta p_μ now going in steps of $\frac{2\pi}{NL}$. This modified proposal has been shown to work in practice resulting in the computation in the chiral condensate in the large N limit of QCD [16]. Currently, work is close to completion in the computation of the pion decay constant, F_π , in the large limit of QCD [17].

II. OVERLAP FERMIONS COUPLED TO LARGE N GAUGE THEORY

Overlap fermions [18] preserve chiral properties of fermions on the lattice and enables a study of fermionic observables in the massless limit of quarks. This makes it possible to compute the low energy parameters in the chiral Lagrangian of large N QCD. In addition, large N gauge fields on the lattice naturally come in disconnected subspaces where each subspace is assigned a topological charge defined using overlap fermions.

The main point is the presence of a lattice phase transition in the large N limit of pure gauge theory which is an extension of the Gross-Witten phase transition [20] in $d = 2$. The single plaquette Wilson action is given by

$$S = bN \sum_{x, \mu \neq \nu} \text{Re Tr} U_{\mu\nu}(x) \quad (1)$$

$$U_{\mu\nu}(x) = U_\mu(x)U_\nu(x + \hat{\mu})U_\mu^\dagger(x + \hat{\nu})U_\nu^\dagger(x) \quad (2)$$

where $b = \frac{1}{g^2 N}$ is kept fixed on the lattice as $N \rightarrow \infty$. The continuum limit corresponds to $b \rightarrow \infty$. The lattice is taken to be a periodic torus with L sites in all directions. The link variables, $U_\mu(x)$, and the plaquette variables, $U_{\mu\nu}(x)$, are in $\text{SU}(N)$. Let $e^{i\theta_{\mu\nu}^k(x)}$; $k = 1, \dots, N$ with $-\pi < \theta_{\mu\nu}^k(x) \leq \pi$ denote the N gauge invariant eigenvalues of $U_{\mu\nu}(x)$. Consider the observable,

$$p(\theta) = \frac{1}{Z} \int [dU_\mu(x)] \frac{1}{NL^d d(d-1)} \sum_{x, \mu \neq \nu, k} \delta[\theta - \theta_{\mu\nu}^k(x)] e^S; \quad Z = \int [dU_\mu(x)] e^S, \quad (3)$$

interpreted as the distribution of the plaquette eigenvalues. This distribution should be uniform at $b = 0$ and should peak around $\theta = 0$ in the weak coupling limit. In the weak coupling limit at any finite N , one expects a sharply peaked distribution at $\theta = 0$ with $p(\pm\pi)$ being highly suppressed. This becomes a phase transition in the $N \rightarrow \infty$ limit at $b = b_c^B$. The distribution is non-zero over the whole range of θ if $b < b_c^B$ and the distribution has a gap with $p(\theta) = 0$ for $|\theta| > \theta_c^B(b)$ if $b > b_c^B$. The critical coupling is exactly at $b_c(B)$ for large L and this is a lattice transition with no continuum counterpart. The phase transition is at $b_c^B = 0.5$ in $d = 2$ for all $L \geq 1$ and is the Gross-Witten phase transition [20]. $b_c^B = 0.4$ in $d = 3$ for all $L \geq 3$ in $d = 3$ [14] and $b_c^B = 0.36$ in $d = 4$ for all $L \geq 9$ in $d = 4$ [15]. It is a very strong first order phase transition in $d = 4$ making it possible to stay in the metastable weak coupling side of the phase for $5 \leq L < 9$ [15].

Gauge fields on the weak coupling side of the above lattice phase transition naturally fall into disconnected pieces. Since there is an upper bound on the plaquette angle, θ , there exist two different gauge fields, U and V say, such that it is not possible to continuously deform U into V without violating the upper bound somewhere in-between. One can conjecture that the disconnected pieces of the gauge field space as dictated by the upper bound on the plaquette angle are in one-to-one correspondence with the topological charge of the gauge field defined using overlap fermions.

Consider a typical update of a large N gauge field on the lattice using a combination of Cabibo-Marinari heat-bath and $SU(N)$ over-relaxation [15] in the weak coupling phase. A cold start where all gauge fields are set to unity will result in thermalized gauge field configurations with zero topological charge. There will be no tunneling events between different topological charges due to the gap in the eigenvalue distribution of the single plaquette. One can obtain thermalized configurations with different topological charges by starting from a gauge field configuration with the appropriate topological charge which also has a gap in the single plaquette eigenvalue distribution. For example, one could start with a “uniform” instanton configuration in $d = 4$. The gauge field is abelian in nature with a uniform field strength equal to $\frac{2\pi}{L^2}$ in a certain plane and a certain direction in color space [16]. Fig. 1 shows thermalized configurations in the $Q = 0$ and $Q = 1$ sectors in the metastable weak coupling phase on a 7^4 lattice with $N = 23$ at $b = 0.35$. The two histories correspond to two different starts where one is a cold start with $Q = 0$ and another is a uniform instanton start with $Q = 1$. No tunneling was observed in either case. The left

panel shows the history of the average plaquette and the right panel shows the distribution of the plaquette eigenvalues. There is a gap in the distribution and both of them look identical. It is useful to note that a naive discretization of the classical $SU(2)$ instanton will not work since its plaquette eigenvalue distribution will not satisfy the gap condition. In practice, the energy distributions in the different topological sectors are very close to each other. Therefore, it is possible to offer one thermalized configuration in one topological




FIG. 1: Plaquette history (left panel) and plaquette eigenvalue distribution (right panel) in the weak coupling phase. Two histories, one thermalized with $Q = 0$ and another with $Q = 1$ are shown at $L = 7$, $N = 23$ and $b = 0.35$.

The gap in the eigenvalue distribution of the plaquette has implications for the overlap Dirac operator in the weak coupling phase of large N QCD. The massless overlap Dirac

operator [19] is defined as

$$D_o = \frac{1 + \gamma_5 \text{sign}(H_w(M))}{2} \quad (4)$$

$H_w(M)$ is the Wilson Dirac operator at mass M , which should be chosen with $-2 < M < 0$ and is an irrelevant parameter in the continuum limit of the theory. Naively, one expects a gap in the spectrum of $H_w(M)$ around zero unless M is close to the critical Wilson coupling and by definition one needs to use a value of M below this critical coupling above. The flow of eigenvalues of $H_w(M)$ as a function of M for a fixed gauge field configuration is directly related to gauge field topology: The net number of levels that cross zero equals the topological charge [21]. One can prove that a gap in the spectrum of the eigenvalue distribution of the plaquette results in the gap in the spectrum of $H_w(M)$ as long as one is significantly away from the critical coupling [22]. Unfortunately, the eigenvalue distribution of the plaquette does not have a gap for $N = 2$ or $N = 3$ when gauge fields are generated using the single plaquette Wilson action. As such, the spectrum of $H_w(M)$ also does not have a gap below the critical coupling [23]. But this is not the case in large N QCD and this gives an unambiguous definition of the topological charge. Furthermore, it also simplifies the numerical evaluation of $\text{sign}(H_w(M))$ which is the most time consuming part in a computation involving the overlap Dirac operator.

III. PHASES OF THE LARGE N GAUGE THEORY

As discussed in the previous section, one has to be in the weak coupling phase of large N QCD on the lattice, defined by the presence of a gap in the eigenvalue distribution of the plaquette, in order to properly reproduce continuum large N QCD. The location of this lattice transition is at $b = b_c^B$. In addition, one has to be in a phase where none of the Z_N symmetries associated with the Polyakov loops in the d directions are broken. We will refer to the strong coupling phase as 0h since it is a “hot” phase with all d Z_N symmetries intact. On the weak coupling side, there will in general be $d + 1$ phases referred to as 0c, 1c, $\dots$, dc where $0, 1, \dots, d$ of the d Z_N symmetries are broken. All the phase exist as long as $L \geq 3$ in $d=3$ and $L \geq 9$ in $d = 4$. Furthermore, one can be in a metastable 0c phase for $5 \leq L < 9$ since the lattice transition separating the 0h and 0c phase at $b = b_c^B$ is strongly first order.

Eguchi-Kawai reduction holds in the 0c phase and large N QCD is properly reproduced in this phase without any finite volume effects. In particular, consider a $K \times K$ Wilson

loop on an L^d periodic lattice with $K \leq L$ or $K > L$. When $K > L$, the Wilson loop is folded since some links appear more than once in the operator. Reduction implies that the eigenvalue distribution of all Wilson loops are independent of L . This is illustrated [14] in Fig. 2 using a 4×4 Wilson loop on 4^3 and 6^3 lattice at a fixed coupling and fixed N . A




FIG. 2: Eigenvalue density distribution of a 4×4 Wilson loop on 4^3 (folded) and 6^3 (unfolded) at $b = 0.66$ and $N = 23$.

proper realization of continuum large N QCD also implies proper scaling of observables in the $0c$ phase and this is illustrated [14] in Fig. 3 using a $L \times L$ Wilson loop on an L^3 lattice for fixed L/b and two different L values. ‘

Unlike the $0h$ to $0c$ transition, the rest of the transitions are physical and therefore, the transition point will scale properly. Our numerical analysis [14, 15] indicates that all




FIG. 3: Eigenvalue density distribution of $L \times L$ Wilson loop on L^3 for $L/b = 5$ and $L = 4, 6$. N is set to 23.

transitions are distinct in the continuum limit but the main focus so far has been on the 0c to 1c transition. One useful observable to locate the phase transition is [8]

$$p(\tilde{P}_\mu) = \frac{1}{N^2} \left\langle \sum_{i,j=1}^N \sin^2 \frac{1}{2} (\theta_i^{\tilde{P}} - \theta_j^{\tilde{P}})^2 \right\rangle \quad (5)$$

where $\tilde{P}_\mu$ is the Polyakov loop (not its trace) in the μ direction located at a fixed point in the 3-plane perpendicular to μ . The $e^{i\theta_j^{\tilde{P}}}$; $j = 1, \dots, N$ are the N eigenvalues of $\tilde{P}_\mu$. The averaging is over all points in the 3-plane and over configurations. Equally spaced eigenvalues respect the $Z(N)$ symmetry in the μ direction and maximize p to 0.5. When the eigenvalue spectrum starts getting modulated and opens a gap, p drops below 0.5. Fig. 4

shows that $b = 0.351$ on 6^4 lattice is in the 0c phase whereas Fig. 5 shows that $b = 0.3568$ on 7^4 lattice is in the 1c phase.




FIG. 4: History of the variable $p(\tilde{P}_\mu)$ for each direction. We see the evolution from a state where one of the four $Z(N)$ factors is broken to one in which all four are preserved.

Let b_c be the location of the numerically obtained transition on an L^d lattice. This can be inverted to define a critical size, $L_c(b)$, that denotes the 0c to 1c transition. The coupling b has dimensions of length in $d=3$ and a numerical analysis [16] shows that $L_c(b) = 4.6(4)b$. The numerically obtained [15] critical size $L_c(b)$ in $d = 4$ shown in Fig. 6 scales according

L=7 N=31 b=0.3568 started from 0c

(measurements taken every 50 lattice passes)




FIG. 5: History of the variable $p(\tilde{P}_\mu)$ for each direction. We see the evolution from a state where all four $Z(N)$ factors are preserved to one where one factor is broken. During the first fifty passes (before the first measurement) Polyakov loops in direction 3 have acquired some structure but, ultimately, direction 2 is selected for breakdown and the Polyakov loops in the other three directions converge to a symmetric state.

to two loop tadpole improved [24] improved perturbation theory as

$$L_c(b) = 0.260(15) \left(\frac{11}{48\pi^2 b e(b)} \right)^{\frac{51}{121}} \exp \left[\frac{24\pi^2 b e(b)}{11} \right] \quad (6)$$

with

$$e(b) = \frac{1}{N} \langle \text{Tr} U_{\mu\nu}(x) \rangle \quad (7)$$




FIG. 6: The transition ranges compared to possible two loop renormalization group curves with tadpole improvement.

The 0c phase of the large N continuum theory on an l^4 torus is clear and it describes large N QCD at zero temperature. The 4c phase can be interpreted as a continuum theory in finite volume and temperature with temperature equal to $1/l$. Rotational invariance is spontaneously broken in the 1c, 2c and 3c phases of the continuum theory. The physical interpretation of these phases is not obvious. The 1c quite possibly describes large N QCD in the deconfined phase since the finite size effects are only felt in the one broken direction and this could be interpreted as finite temperature. Then, one would conclude that large N QCD in the confined phase does not depend on temperature, a view that has been recently discussed [25]. The 3c phase possibly describes a continuum theory in a small finite box at zero temperature. The continuum theories described in the phases other than 0c and 1c might have implications not directly related to large N QCD. A careful study of the 0c to 1c transition would tell us if the transition is the physical finite temperature phase transition. The critical box size [15] associated with the 0c to 1c transition in units of string tension is $l_c\sqrt{\sigma} \approx 1.56$. A comparison with the numerical estimates of the critical temperature [26] indicate that $l_c T_c = 1$. Recent numerical studies indicate a finite latent heat associated with this transition [27] providing further supporting evidence for this to be the physical finite temperature phase transition.

IV. SPONTANEOUS CHIRAL SYMMETRY BREAKING

Spontaneous chiral symmetry breaking is an important phenomenon in QCD. The mass of the pion made up of two light quarks is not equal or close to the sum of quark masses but proportional to the square root of the quark mass since the pion is a pseudo-Goldstone boson associated with spontaneous chiral symmetry breaking in QCD with massless quarks. Several lattice studies have investigated this phenomenon in the past using Wilson fermions [28] and staggered fermions [29] and more recently using overlap fermions [30]. Quenched lattice studies of this phenomenon suffer from quenched pathologies. The quenched pathologies arise out of the fact that the singlet meson, η' , becomes a pseudo-Goldstone boson in the quenched approximation [31]. This introduces a new mass parameter m_0 in the chiral Lagrangian with $m_0 \approx 900\text{MeV}$ in the chiral limit. The pion mass no longer is linear

in the square root of the light quark mass and instead is given by [31]

$$m_\pi^2 \propto m^{\frac{1}{1+\delta}}; \quad \delta = \frac{m_0^2}{N(4\pi f_\pi)^2} \approx \frac{0.6}{N}. \quad (8)$$

where m is the light quark mass and $f_\pi = 93\text{MeV}$ is the pion decay constant. Spontaneous chiral symmetry breaking occurs due to the presence of a non-zero density of eigenvalues of the massless Dirac operator near zero. Let $\rho(\lambda)$ denote the density of the paired non-zero $\pm i\lambda$ of the anti-hermitian massless Dirac operator after removing the exact zero eigenvalues due to gauge field topology. The chiral condensate [33],

$$\lim_{m \rightarrow 0} \lim_{V \rightarrow \infty} \langle \bar{\psi}\psi(m) \rangle = \pi\rho(0), \quad (9)$$

is non-zero if $\rho(0)$ is non-zero and this shows spontaneous chiral symmetry breaking. The effect of the m_0 term in quenched QCD results in a diverging chiral condensate [31], given by

$$\lim_{V \rightarrow \infty} \langle \bar{\psi}\psi(m) \rangle \propto m^{-\frac{\delta}{1+\delta}} \quad (10)$$

Typical quenched calculations on the lattice focus on the behavior of pion mass as a function of quark mass to extract the quenched divergence. Since $\delta > 0$, the effect of quenched divergence is to make the pion mass at a fixed quark mass a little heavier than what it should be in unquenched QCD. Since all calculations are performed at finite volume, the effect due to quenched divergence cannot be distinguished from finite volume effects [32]. On the other hand, the chiral condensate diverges in the quenched approximation and this effect goes in the opposite direction from that of finite volume effects. In the large N limit, δ goes to zero and there are no quenched pathologies.

In order to avoid quenched pathologies, one has to take the large N limit before one takes the infinite volume limit. Furthermore, Eguchi-Kawai reduction holds in the 0c phase and therefore the infinite N limit in this phase will not depend on the physical volume. We should therefore see evidence for a chiral condensate on a finite lattice in the large N limit as long as we are in the 0c phase. This is counter-intuitive since we expect to see a finite density of eigenvalues near zero in the spectrum of the massless Dirac operator even at finite lattice volume. Chiral random matrix theory [34] provides an understanding of spontaneous chiral symmetry breaking in large N QCD on a finite lattice [16].

Simple counting of N degrees of freedom shows that $\langle \bar{\psi}\psi \rangle$ is proportional to N . Therefore the low lying eigenvalues of the massless Dirac operator should scale like $\frac{1}{NV}$ if chiral symmetry is spontaneously broken in the large N limit. This is a consequence of level repulsion.

The paired non-zero eigenvalues, $\pm\lambda$, of the massless hermitian overlap Dirac operator satisfies $0 < \lambda < 1$. The operator has $2NV$ positive eigenvalues (assuming zero topology) on a $V = L^4$ lattice. Level repulsion would result in eigenvalues that are roughly equally spaced. Therefore, the mean spacing will be proportional to $\frac{1}{NV}$ and the lowest eigenvalues will scale like $\frac{1}{NV}$. Furthermore, the scaled eigenvalues $z = \lambda\Sigma NV$ should essentially be distributed in some universal manner and this universal behavior is dictated by chiral random matrix theory [35]. The scale, Σ , that connects z and λ is the chiral condensate. Universal behavior is approached as NV , the number of eigenvalues, go to infinity. This can be achieved by keeping N fixed and taking V to infinity which is the conventional approach. But, it can also be achieved by taking N to infinity at fixed V and this makes use of the reduction in large N QCD. Quenched divergences imply that λ does not scale like $\frac{1}{NV}$ at fixed N as $V \rightarrow \infty$. In spite of this, it has been shown that one can define a scale $\Sigma(V)$ such that chiral random matrix theory is respected in the quenched approximation [36]. But, $\Sigma(V)$ diverges as $V \rightarrow \infty$.

One could show evidence for spontaneous chiral symmetry breaking in large N QCD by a direct computation of Σ in the 0c phase. This can be done by a computation of $\lim_{m \rightarrow 0} \lim_{N \rightarrow \infty} \langle \bar{\psi}\psi(m) \rangle$ on a finite L^d lattice. In this case, it is necessary to take the large N limit before taking the massless limit since the chiral condensate is zero at $m = 0$ for any finite N . Numerically, one has to perform some sort of a finite N analysis equivalent to the usual finite volume analysis done in order to extract an order parameter in the thermodynamic limit. Chiral random matrix theory provides us with a way to perform the finite N analysis without a computation of the chiral condensate. Let λ_i , $i = 1, \dots, 2NV$, with $\lambda_1 < \lambda_2 < \dots < \lambda_{2NV}$ be all the positive eigenvalues of the hermitian overlap Dirac operator. Let $z_k = \lambda_k \Sigma NV$ be the scaled eigenvalues that obey the universal distribution given by chiral random matrix theory [37]. As N increases, more and more z_k will obey the universal distributions. In particular, one will find that $p(r = \frac{\lambda_1}{\lambda_2})$ obeys a universal distribution as long as N is large enough. Once this is achieved, Σ can be obtained by matching $\langle z_1 \rangle$ and $\langle z_2 \rangle$ to their values given by chiral random matrix theory. A numerical computation shows that one can obtain Σ in the large N limit of QCD using chiral random matrix theory [16]. Existence of a chiral condensate in the continuum limit amounts to showing that $\Sigma(b)$ obtained on the lattice scales properly in the continuum limit and one finds that $\Sigma^{1/3} l_c = 0.588$ which translates to $\Sigma^{1/3} \approx 155 \text{ MeV}$ if we assume that $\frac{1}{l_c} = T_c = 264 \text{ MeV}$.

A. Chiral condensate in the large N limit of QCD_2

Before proceeding to the computation of chiral condensate using chiral random matrix theory, it is useful to verify the ideas of chiral random matrix theory in two dimensional QCD [16]. The chiral condensate in the 't Hooft model is exactly known [38] and it is given by

$$\Sigma(b) = \frac{|M|}{\sqrt{6\pi b}} \quad (11)$$

for the overlap Dirac operator, where M is the Wilson mass parameter used in the definition of the overlap Dirac operator. Eguchi-Kawai reduction holds for all L in $d = 2$ and therefore one can compute the chiral condensate by taking the large N limit on a 1^2 lattice. One can compute the full spectrum of the overlap Dirac operator for very large N using exact diagonalization on a 1^2 lattice. Let $p_1(z_1)$ and $p_2(z_2)$ be the universal distributions of the two lowest eigenvalues and let $p(r)$ be the distribution of the ratio of the first eigenvalue and the second eigenvalue. Fig. 7 shows that approach to chiral random matrix theory as N is increased on a 1^2 lattice at a fixed b . $N = 37$ is too small for chiral random matrix theory to be satisfied. At $N = 79$ the first eigenvalue already satisfies the universal distribution and both eigenvalues satisfy the universal distribution beyond $N = 137$. The coupling was set to $b = 1$ and we used $M = -1$ in the definition of the overlap Dirac operator. Therefore, we should find $\Sigma = 0.23$ as $N \rightarrow \infty$. Furthermore, $\langle r \rangle = 0.37$ as $N \rightarrow \infty$. The approach to the $N \rightarrow \infty$ limit of $\langle r \rangle$ and Σ using the first and second eigenvalue is shown in Fig. 8. One can see that the first eigenvalue approaches chiral random matrix theory before the second eigenvalue.

B. Chiral condensate in the large N limit of QCD

Analysis of QCD_2 shows that one can use chiral random matrix theory to extract the chiral condensate in the large N limit of QCD. To do this, one has to be in the $0c$ phase of QCD. Therefore, we worked with L in the range of 6 to 10. At each L , there is a range of coupling $b_c^B < b < b_c(L)$ where the theory is in the $0c$ phase. It is best to pick the coupling close to $b_c(L)$ to minimize lattice spacing effects at that L . We therefore used [15] $b = 0.350$ for $L \geq 6$, $b = 0.355$ for $L \geq 7$ and $b = 0.3585$ for $L \geq 8$. We worked with $L = 6, 7, 8$ at $b = 0.350$ to ensure that reduction holds. We worked with $L = 8, 9, 10$ at $b = 0.355$ again




FIG. 7: Approach to chiral random matrix theory on a 1^2 lattice at $b = 1$ as N is increased from 37 to 273.




FIG. 8: Estimation of the chiral condensate in the $N \rightarrow \infty$ limit of QCD_2 at $b = 1$ as N is increased from 23 to 273.

to ensure that reduction holds. We worked with $L = 9$ at $b = 0.3585$ and we also obtained one result on a coarse lattice of $b = 0.346$ at $L = 9$. The scaling properties of the code are as follows. The gauge field generation scales like $N^3 L^4$. The computation of the lowest two eigenvalues of the hermitian overlap Dirac operator scales like $N^3 L^8$ and this arises from

two factors. One factor is due to the action of the Wilson Dirac operator on a vector and this scales as $N^2 L^4$. The low lying eigenvalues of the overlap Dirac operator are obtained using Ritz techniques and the relevant condition number is the ratio of the highest to the lowest eigenvalue and therefore the second factor scales as $N L^4$.

Fig. 9 shows the approach to chiral random matrix theory as one approaches the large N limit at a fixed L and b . One sees a clear agreement with chiral random matrix theory in the distribution of the ratio of the two lowest eigenvalues for $N \geq 23$. All configurations are in the zero topological sector. The universal distribution of $p(r)$ is different in the different topological sectors and Fig. 10 shows that one finds agreement with chiral random matrix theory in the $Q = 1$ topological sector when there is agreement in the $Q = 0$ topological sector. Finally, Fig. 11 shows that one also finds agreement with chiral random matrix theory as one goes toward the continuum limit.

The first and second eigenvalues are used to obtain the chiral condensate, Σ , by setting the average of $z_i = \lambda_i \Sigma_i N L^4$; $i = 1, 2$ equal to the ones dictated by chiral random matrix theory. One should find $\Sigma_1 = \Sigma_2$ for values of b , L and N where $p(r)$ agreed with the universal distribution given by chiral random matrix theory. Fig. 12 shows the results for Σ_1 and Σ_2 from all our numerical simulations at $b = 0.350$ where we found $p(r)$ to agree with chiral random matrix theory. First of all we find that $\Sigma_1 = \Sigma_2$ for all the data shown in Fig. 12. Furthermore, we do not see any finite volume effects at $b = 0.350$ indicating that reduction works properly for fermionic observables. The finite N effects are relatively small over the range of N shown since the y -axis in Fig. 12 shows a deviation of only 5%. We can therefore estimate $\Sigma^{1/3}(b = 0.350) = 0.142 \pm 0.006$.

Estimation of the chiral condensate using chiral random matrix theory has proved to be far superior to conventional techniques [16]. The main problem that plagues the conventional approach is the quadratic divergence in $\langle \bar{\psi} \psi \rangle$. Therefore, large subtractions are needed in order to extract the chiral condensate. This can be remedied by considering derivatives with respect to m^2 of the chiral condensate. This ameliorates the problem but the dependence on small quark masses was still strong and this makes it difficult to estimate the chiral condensate using extrapolation [16]. In addition to the need for subtractions, one can only obtain a stochastic estimate of this quantity and it is necessary for the random source to have good overlap with the low lying eigenvalues. On the other hand, numerical evaluation using Ritz functional techniques results in estimates of low lying eigenvalues with errors that




FIG. 9: Approach to chiral random matrix theory at a fixed b and L with increasing N .

are less than a tenth of a percent.

If we interpret the 0c to 1c transition as a finite temperature phase transition, then one would expect chiral symmetry to be restored in the 1c phase. But, chiral random matrix theory arguments would indicate that chiral symmetry would still be broken since the Z_N




FIG. 10: Evidence for agreement with chiral random matrix theory in the $Q = 0$ and $Q = 1$ topological sectors at a fixed $b = 0.350$.




FIG. 11: Evidence for agreement with chiral random matrix theory as one approaches the continuum limit.




FIG. 12: Estimate of the chiral condensate in the large N limit of QCD.

symmetry is broken only in one of the four directions. Three other directions still behave as if they were infinite in extent and one would still expect level repulsion with level spacings that go like $\frac{1}{NL^3}$. Therefore, one would expect chiral symmetry to be broken in the 1c phase with a value that depends on the length of the direction which is broken. Therefore, it would be interesting to compute the chiral condensate in the 1c phase. Similar arguments

would also say that chiral symmetry is broken in the 2c and 3c phase. Chiral symmetry is expected to be restored only in the 4c phase. A study of chiral symmetry breaking in these other phases would be interesting on their own right for a full understanding of the associated continuum theories.

V. CONCLUSIONS AND FUTURE WORK

Phase transitions as a function of scale are ubiquitous in large N QCD. Continuum reduction [14] says that large N gauge theory on an l^d torus undergoes a phase transition when $l = l_c$ and large N QCD is properly reproduced as long as $l > l_c$. This can be used to numerically solve large N QCD by working on small lattices. Furthermore, one can work in the quenched approximation on the lattice. It is important to take the $N \rightarrow \infty$ limit before one takes any other limit. Then, one finds that there are no finite volume effects and it is possible to obtain the low energy parameters of the chiral Lagrangian in the large N limit of QCD [16]. Chiral random matrix theory [34] proved to be a useful tool in the extraction of the chiral condensate since it describes the behavior of small eigenvalues for large but finite number of degrees of freedom.

The next step is the computation of the pion decay constant and work is close to completion [17]. This will establish the validity of momentum quenching in the computation of meson propagators. A study of current correlators will be useful in the extraction of the mass of the ρ .

On the technical side, the Wilson Dirac operator has a gap in its spectrum at values of M used in the overlap Dirac operator. This is due to the lattice phase transition present in the large N limit of QCD and is a consequence of the gap in the eigenvalue distribution of the single plaquette. Therefore, gauge fields come in disconnected subspaces. In addition, the gap in $H_w(M)$ results in a significant reduction in the computational cost of the action of the overlap Dirac operator on a vector.

The 1c to 4c phases discussed in section 3 have possible implications for certain string theories [4] and it would be useful to compute the chiral condensate in all these phases.

Acknowledgments

R. N. acknowledges partial support by the NSF under grant number PHY-0300065 and also partial support from Jefferson Lab. The Thomas Jefferson National Accelerator Facility (Jefferson Lab) is operated by the Southeastern Universities Research Association (SURA) under DOE contract DE-AC05-84ER40150. H. N. acknowledges partial support by the DOE under grant number DE-FG02-01ER41165. A significant part of the results presented here were obtained in collaboration with Joe Kiskis. We thank the organizers of the workshop for providing a stimulating environment.

-
- [1] G. 't Hooft, *Nucl. Phys.* **B72**, 461 (1974); E. Witten, *Nucl. Phys.* **B160**, 57 (1979); S. Coleman, *Lectures at the 1979 International School of Subnuclear Physics, Erice, 1979*.
 - [2] G. 't Hooft, *Nucl. Phys.* **B75**, 461 (1974); in: A. Zichichi (Ed.), *New Phenomena in Subnuclear Physics, Part A, Proceedings of the International School of Subnuclear Physics, Erice, 1975*.
 - [3] A.V. Manohar, *Talk at the Les Houches Summer School in Theoretical Physics, Session 68: Probing the Standard Model of Particle Interactions, Les Houches, France, 28 Jul - 5 Sep 1997*.
 - [4] O. Aharony, S.S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, *Phys. Rept.* **323**, 183 (2000).
 - [5] Contribution by M. Teper to this proceedings.
 - [6] S. Sharpe, in *Lectures at TASI 1994: CP Violation and the Limits of the Standard Model*, World Scientific, Ed. J. Donoghue.
 - [7] T. Eguchi and H. Kawai, *Phys. Rev. Lett.* **48**, 1063 (1982).
 - [8] G. Bhanot, U.M. Heller and H. Neuberger, *Phys. Lett.* **B112**, 47 (1982).
 - [9] A. Gonzalez-Arroyo and M. Okawa, *Phys. Lett.* **B120**, 174 (1983).
 - [10] A. Gonzalez-Arroyo and C.P. Korthals Altes, *Phys. Lett.* **B131**, 396 (1983).
 - [11] D. Gross and Y. Kitazawa, *Nucl. Phys.* **B206**, 440 (1982).
 - [12] H. Levine and H. Neuberger, *Phys. Lett.* **B119**, 183 (1982).
 - [13] J. Kiskis, R. Narayanan and H. Neuberger, *Phys. Rev.* **D66**, 025019 (2002).
 - [14] R. Narayanan and H. Neuberger, *Phys. Rev. Lett.* **91**, 081601 (2003).
 - [15] J. Kiskis, R. Narayanan and H. Neuberger, *Phys. Lett.* **B574**, 65 (2003).

- [16] R. Narayanan and H. Neuberger, *Nucl. Phys.* **B696**, 107 (2004).
- [17] R. Narayanan and H. Neuberger, *Manuscript in preparation*.
- [18] R. Narayanan and H. Neuberger, *Nucl. Phys.* **B443**, 305 (1995).
- [19] H. Neuberger, *Phys. Lett.* **B417**, 141 (1998).
- [20] D. Gross and E. Witten, *Phys. Rev.* **D21**, 446 (1980).
- [21] R. Narayanan and H. Neuberger, *Phys. Rev. Lett.* **71**, 3251 (1993).
- [22] H. Neuberger, *Phys. Rev.* **D61**, 085015 (2000).
- [23] R.G. Edwards, U.M. Heller and R. Narayanan, *Nucl. Phys.* **B535**, 403 (1998).
- [24] P. Lepage, *Lectures at 1996 Schladming Winter School on Perturbative and Nonperturbative Aspects of Quantum Field Theory (Schladming, Austria, March 1996)*, hep-lat/9607076.
- [25] T.D. Cohen, *Phys. Rev.* **D70**, 014011, 2004.
- [26] B. Lucini, M. Teper and U. Wenger, *Phys. Lett.* **B545**, 197 (2002).
- [27] J. Kiskis, *Manuscript in preparation*.
- [28] H. Thacker, W.A. Bardeen, A. Duncan and E. Eichten, *Nucl. Phys. Proc. Suppl.* **73**, 243 (1999); S. Aoki, G. Boyd, R. Burkhalter, S. Ejiri, M. Fukugita, S. Hashimoto, Y. Iwasaki, K. Kanaya, T. Kaneko, Y. Kuramashi, K. Nagai, M. Okawa, H. P. Shanahan, A. Ukawa and T. Yoshie, *Nucl. Phys. Proc. Suppl.* **73**, 189 (1999)
- [29] S. Kim and D.K. Sinclair, *Phys. Rev.* **D52**, R2614 (1995); S. Kim and S. Ohta, *Phys. Rev.* **D61**, 074506 (2000).
- [30] T-W. Chiu and T-H. Hsieh, *Phys. Rev.* **D66**, 014506 (2002); Y. Chen, S.J. Dong, T. Draper, I. Horvath, F.X. Lee, K.F. Liu, N. Mathur and J.B. Zhang, *Phys. Rev.* **D70**, 034502 (2004).
- [31] S. Sharpe, *Phys. Rev.* **D46**, 3146 (1992).
- [32] R.D. Mawhinney, *Nucl. Phys. Proc. Suppl.* **47**, 557 (1996).
- [33] T. Banks and A. Casher, *Nucl. Phys.* **B169**, 103 (1980).
- [34] J.J.M. Verbaarschot and T. Wettig, *Annu. Rev. Nucl. Part. Sci.* **50**, 343 (2000).
- [35] E.V. Shuryak and J.J.M. Verbaarschot, *Nucl. Phys.* **A560**, 306 (1993).
- [36] P.H. Damgaard, *Nucl. Phys.* **B608**, 162 (2001).
- [37] P.H. Damgaard and S.M. Nishigaki, *Phys. Rev.* **D63**, (2001) 045012; hep-th/0006111.
- [38] A.R. Zhitnitsky, *Phys. Lett.* **B165**, 405 (1985).